



$$E = \frac{\sqrt{+ -}}{-} = - -$$

2.2.1. *S a ial āng la i ʒ of*

$$- = \frac{1}{\sqrt{}} + \frac{1}{\sqrt{}} - \frac{1}{\sqrt{}}$$

a finite resolution then either the physics or numerics define a finest length scale (Δx) and we can pick $\Delta x \gg \lambda^{-2}$

3.3. Application

Finally, the expansions were tested for various hydrogen-like sys-

- [10] M. Reiher, Relativistic Douglas-Kroll-Hess theory, *Wiley Interdiscip. Rev.: Comput. Mol. Sci.* 2 (2012) 139–149.
- [11] K.G. Dyall, An exact separation of the spin-free and spin-dependent terms of the Dirac-Coulomb-Breit Hamiltonian, *J. Chem. Phys.* 100 (3) (1994) 2118–2127.
- [12] K.G. Dyall, Interfacing relativistic and nonrelativistic methods. I. Normalized elimination of the small component in the modified Dirac equation, *J. Chem. Phys.* 106 (23) (1997) 9618–9626.
- [13] L. Foldy, S. Wouthuysen, On the Dirac theory of spin 1/2 particles and its non-relativistic limit, *Phys. Rev.* 78 (1) (1950) 29–36.
- [14] C. van Wullen, On the eigenfunctions of the Douglas-Kroll operator, *Chem. Phys.* 356 (2009) 199–204.
- [15] B. Alpert, G. Beylkin, D. Gines, L. Vozovoi, Adaptive solution of partial differential equations in multiwavelet bases, *J. Comput. Phys.* 182 (1) (2002) 149–190.
- [16] R. Harrison, G. Fann, T. Yanai, G. Beylkin, Multiresolution quantum chemistry in multiwavelet bases, in: P.M.A. Sloot, et al. (Ed.), *Lecture Notes in Computer Science, Computational Science-ICCS 2003*, vol. 2660, Springer, 2003, pp. 103–110.
- [17]