

1. [2360/030922 (10 pts)] Given the matrices

$$A = \begin{pmatrix} 2 & 1 & 3 \\ 4 & 3 & 5 \\ 1 & 2 & 1 \end{pmatrix} \quad B = \begin{pmatrix} 2 & 1 & 3 \\ 0 & 1 & 2 \end{pmatrix} \quad C = \begin{pmatrix} 1 & 4 \\ 1 & 1 \end{pmatrix}$$

write the word TRUE or FALSE as appropriate. No work need be shown, no work will be graded and no partial credit will be given.

- (a) $CB = \begin{pmatrix} 2 & 3 \\ 4 & 5 \\ 1 & 1 \end{pmatrix}$ (b) $\text{Tr } B^T A^T = 2$ (c) $A^T A = A A^T$ (d) $\sum_j C^T C = 10$ (e) $AB - A^T B^T$ is not defined

SOLUTION:

- (a) **FALSE** $CB = \begin{pmatrix} 1 & 4 & 2 & 1 & 3 \\ 0 & 1 & 2 \\ 0 & 2 & 3 \end{pmatrix} = \begin{pmatrix} 2 & 5 & 11 \\ 1 & 2 \\ 2 & 6 & 2 \end{pmatrix}$
- (b) **TRUE** $\text{Tr } B^T A^T = \text{Tr } A B^T = \text{Tr } \begin{pmatrix} 2 & 0 & 3 \\ 1 & 15 & 1 \\ 3 & 2 & 0 \end{pmatrix} = \text{Tr } \begin{pmatrix} 2 & 6 & 2 \\ 1 & 1 & 15 \\ 3 & 1 & 1 \end{pmatrix} = 2 + 1 + 1 = 2$
- (c) **FALSE** $A^T A$ is $(2 \ 3)(3 \ 2) = 2 \ 2$ whereas $A A^T$ is $(3 \ 2)(2 \ 3) = 3 \ 3$ so they cannot be equal
- (d) **FALSE** $\sum_j C^T C = \begin{pmatrix} 1 & 1 & 4 & 3 \\ 4 & 1 & 4 & 3 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 4 \\ 4 & 16 \end{pmatrix} \neq A A^T$

We need to find constants $c_1; c_2; c_3$ such that $c_1 \neq$

(b)

$$\begin{aligned}
 A^T A \vec{x} &= \begin{pmatrix} 2 & 3 \\ 1 & 2 \\ 4 & 5 \\ 1 & 1 \end{pmatrix} \vec{x} \\
 A^{-1} A^T A \vec{x} &= A^{-1} \begin{pmatrix} 2 & 3 \\ 1 & 2 \\ 4 & 5 \\ 1 & 1 \end{pmatrix} \vec{x} \quad \text{Note: } A^{-1} A^T = I \text{ and } I A = A \\
 A \vec{x} &= A^T \begin{pmatrix} 2 & 3 \\ 1 & 2 \\ 4 & 5 \\ 1 & 1 \end{pmatrix} \vec{x} \\
 A^{-1} A \vec{x} &= A^{-1} A^T \begin{pmatrix} 2 & 3 \\ 1 & 2 \\ 4 & 5 \\ 1 & 1 \end{pmatrix} \vec{x} \quad \text{Note: } A^{-1} A = I \text{ and } I \vec{x} = \vec{x} \\
 \vec{x} &= A^{-1} A^{-1 T} \begin{pmatrix} 2 & 3 \\ 1 & 2 \\ 4 & 5 \\ 1 & 1 \end{pmatrix} \vec{x} \quad \text{Note: } A^{-1 T} = (A^{-1})^T \\
 \vec{x} &= \begin{pmatrix} 2 & 3 & 2 & 3 \\ 1 & 0 & 1 & 0 \\ 4 & 0 & 1 & 0 \\ 3 & 0 & 2 & 1 \end{pmatrix} \begin{pmatrix} 2 \\ 1 \\ 4 \\ 1 \end{pmatrix} \\
 \vec{x} &= \begin{pmatrix} 2 & 3 & 2 & 3 \\ 2 & 0 & 5 & 1 \\ 4 & 0 & 1 & 0 \\ 5 & 0 & 13 & 1 \end{pmatrix} \begin{pmatrix} 2 \\ 1 \\ 4 \\ 1 \end{pmatrix} = \begin{pmatrix} 32 \\ 7 \\ 4 \\ 18 \end{pmatrix}
 \end{aligned}$$

5. [2360/030922 (12 pts)] Determine if each of the following sets of vectors forms a basis for \mathbb{R}^3 . Justify your answers.

$$\begin{aligned}
 \text{(a)} \quad & \begin{pmatrix} 8 & 2 & 3 \\ 2 & 1 & 3 \\ 4 & 2 & 5 \\ 0 & 1 & 1 \end{pmatrix} \begin{pmatrix} 2 \\ 3 \\ 15 \\ 1 \end{pmatrix} \\
 \text{(b)} \quad & \begin{pmatrix} 8 & 2 & 3 \\ 2 & 1 & 3 \\ 4 & 2 & 5 \\ 0 & 1 & 2 \end{pmatrix} \begin{pmatrix} 2 \\ 3 \\ 15 \\ 8 \\ 2 \end{pmatrix}
 \end{aligned}$$

SOLUTION:

Note that the dimension of \mathbb{R}^3 is 3 so a basis consists of 3 linearly independent vectors.

- (a) The set contains only 2 vectors and thus cannot form a basis for \mathbb{R}^3 regardless of the linear dependence or independence of the vectors in the set.
- (b) Three vectors in \mathbb{R}^3 can potentially be a basis if they are linearly independent. To check for this, we need to see if the only solution to

$$c_1 \begin{pmatrix} 2 & 3 \\ 1 & 0 \\ 4 & 2 \\ 0 & 1 \end{pmatrix} + c_2 \begin{pmatrix} 2 & 3 \\ 3 & 1 \\ 4 & 15 \\ 1 & 1 \end{pmatrix} + c_3 \begin{pmatrix} 2 & 3 \\ 3 & 0 \\ 4 & 8 \\ 2 & 0 \end{pmatrix} = \begin{pmatrix} 2 \\ 1 \\ 4 \\ 0 \end{pmatrix} \Rightarrow \begin{pmatrix} 2 & 3 & 3 & 2 \\ 1 & 3 & 0 & 1 \\ 4 & 1 & 8 & 4 \\ 0 & 1 & 2 & 0 \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \\ c_3 \\ 0 \end{pmatrix} = \begin{pmatrix} 2 \\ 1 \\ 4 \\ 0 \end{pmatrix}$$

is the trivial solution. The determinant of the coefficient matrix is

$$\begin{vmatrix} 1 & 3 & 3 \\ 2 & 1 & 8 \\ 0 & 1 & 2 \end{vmatrix} = 1(1-6) + 2(3-2) = 1(-5) + 2(1) = -3 \neq 0$$

implying that the system has nontrivial solutions, further implying that the vectors are linearly dependent and thus cannot form a basis for \mathbb{R}^3 .

6. [2360/030922 (24 pts)] The following parts are unrelated.

$$\text{(a) (12 pts) Find the RREF of } A = \begin{pmatrix} 2 & 3 & 1 & 9 \\ 4 & 1 & 1 & 15 \\ 3 & 11 & 5 & 35 \end{pmatrix}$$

(b) Let $\vec{v} = \begin{pmatrix} 2 \\ p+q \\ r \\ s \end{pmatrix} \in W$ with $s > 0$. Then $1\vec{v} = \begin{pmatrix} 2 \\ p+q \\ r \\ s \end{pmatrix} \notin W$ since $s < 0$. This implies that W is not closed under scalar multiplication and thus is not a subspace.