

DECONVOLUTION AND INVERSION

Proceedings of a
workshop sponsored by the
European Association of Exploration Geophysicists,
the Society of Exploration Geophysicists

the National Research Council of Italy and the
National Science Foundation in the United States,
Rome, 3-5 September 1986

ORGANIZED BY
M. BERNABINI
P. CARRION, G. JACOVITTI
F. ROCCA S. TRETEI

MATHEMATICAL THEORY FOR SEISMIC

G. BEYLKIN

INTRODUCTION

Migration methods in seismics are computationally intensive and powerful tools for interpretation of seismic experiments. A partial list of papers on migration methods in the geophysical literature include those by Hagedoorn (1954), Lindsey and Herman (1970), Rockwell (1971), Claerbout (1971), Schneider (1971, 1978), Claerbout and Doherty (1972), French (1974, 1975), Gardner *et al.* (1974), Cohen and Bleistein (1977, 1979), Stolt (1978), Berkhout (1980, 1984), Clayton and Stolt (1981), Johnson and French (1982), Devaney (1984), Gazdag and Squazzero (1984), Tarantola (1984), Bleistein and Gray (1985), Stolt and Wacklin (1985) and many more. This book

derivation of migration algorithms to solve the linearized inverse scattering problem is best understood within the theory of pseudodifferential and Fourier integral operators.

The sequence for the derivation of a migration algorithm within this theory can be briefly described as follows. For a given background model M_0 and a given set of data D , one

17.1 PSEUDODIFFERENTIAL AND FOURIER INTEGRAL OPERATORS

The title of this section might discourage a non-mathematician since these notions are not yet a part of the standard curriculum in applied mathematics. However, these mathematical objects are well known under different names to electrical engineers.

A more symmetric form of the symbol with respect to α and β is

[REDACTED]

The Sobolev space $H^s(R^d)$ is the space of distribution in R^d whose Fourier transform is a square-integrable function in R^d with the measure $(1 + |k|^2)^s dk$. It is a Hilbert space with the inner product

$$(u, v) = \frac{1}{(2\pi)^d} \int_{R^d} \hat{u}(k) \bar{\hat{v}}(k) (1 + |k|^2)^s dk, \quad (17.11)$$

where the bar denotes complex conjugation.

The subspace $H_{\text{comp}}^s(Q)$ of $H^s(R^d)$ consists of the distributions with the support in the compact set Q ; $H_{\text{comp}}^s(X)$ is the union of the spaces $H_{\text{comp}}^s(Q)$, where Q spans the collection of all compact subsets of X . Finally, $H_{\text{loc}}^s(X)$ is a space of distributions in X , such that if properly localized (by infinitely differentiable cutoff functions to compact subsets exhausting X) these distributions belong to $H_{\text{comp}}^s(Q)$.

operator $\mathbf{P} \in L^m$ an asymptotic expansion of this operator can be constructed:

$$\mathbf{P} = \mathbf{T}_m + \mathbf{T}_{m-1} + \mathbf{T}_{m-2} + \dots, \quad (17.13)$$

where

$$\mathbf{T}_j \in L^j(X), \quad (17.14)$$

for $j = m, m-1, m-2, \dots$, and

$$(\mathbf{P} - \mathbf{T}_m - \mathbf{T}_{m-1} - \dots - \mathbf{T}_j) \in L^{j-1}(X), \quad (17.15)$$

for $j = m, m-1, m-2, \dots$. Such an expansion is modulo regularizing operators

Fourier integral operators

An operator of the form

$$(Ff)(x) = \frac{1}{(2\pi)^n} \int \int f(y) A(x, y, k) e^{i\Phi(x, y, k)} dy dk \quad (17.16)$$

Given the background index of refraction n (background model), the linearized

inverse scattering problem is that of characterization of the perturbation f using observations of the scattered field u on the boundary ∂X of the region X .

If the propagation is governed by the Helmholtz equation, then the scattered field $\hat{u}(s, r, \omega)$ satisfies within the single scattering (or distorted wave Born) approximation the following integral equation

$$\hat{u}(s, r, \omega) = -\omega^2 \int_X G(y, r, \omega) f(y) G(s, y, \omega) dy, \quad (17.19)$$

where G is the Green's function of the background model.

The Green's function G is the solution of the equation

$$(\nabla_y^2 + \omega^2 n_0^2(y)) G(y, r, \omega) = \delta(y - r), \quad (17.20)$$

and, in principle, can be computed given the background model. The incident field

where the function $h(s, r, x)$ is yet to be described. Here Re denotes the real part of the

where

$$\Phi(s, r, x) = \phi(s, x) + \phi(x, r), \quad (17.29)$$

is the total travel time between the source, the point x and r .

Step 2. At this step we localize the computation to the neighbourhood of the point of reconstruction x . If $\varepsilon < |x - y|$, where ε is any positive number, then the result of integration in (17.28) (over the part of the domain X described by this condition) is infinitely differentiable and, therefore, will not affect the asymptotics. If $|x - y| < \varepsilon$ we replace the phase of the exponent by the first term of the Taylor series

$$\Phi(s, r, x) - \Phi(s, r, y) = \nabla_x \Phi(s, r, x) \cdot (x - y), \quad (17.30)$$

and 'freeze' the value of the amplitude terms at the point x . By doing this we account for the most singular term in the asymptotic expansion with respect to smoothness. We obtain from (17.13)

$$\mathcal{L}_{\omega}(x) = \frac{1}{\omega} \operatorname{Re} \int_0^{\infty} \int \int_{\partial X} \frac{e^{i\omega \nabla_x \Phi(s, r, x) \cdot (x - y)} L(s, r, x)}{\dots} \dots$$

Step 3. At this step we set $\omega^2 h(s, r, x)$ to be the Jacobian of the change of variables from $\omega \in [0, \infty]$ and $r \in \partial X_r$ to $k \in R^3$. We have

$$k = \omega \nabla_x \Phi(s, r, x), \quad (17.32)$$

so that

$$dk = h(s, r, x) dr \omega^2 d\omega. \quad (17.33)$$

The function $h(s, r, x)$ can be computed by ray tracing using the identity (Beylkin 1985a)

coverage in the space of spatial frequencies. This domain is determined by the map

we have

$$f_{\text{out}}(s) = -\frac{1}{2\pi} \int \frac{h(s, r, x)}{|r - x|} \dots$$

Concluding remarks

This paper demonstrates (without proofs) how to derive migration algorithms using tools from the theory of pseudodifferential and Fourier integral operators. The purpose of this presentation is to discuss the mathematical technique as it is applied in the context of seismic problems rather than to propose a specific algorithm. For this reason, instead of giving numerical examples, I refer to the papers Miller *et. al.* (1984, 1987), and Beylkin *et. al.* (1985), where specific algorithms are presented along with

- STOLT, R. H. and WEGLEIN, A. B. 1985, Migration and inversion of seismic data, *Geophysics*, 50, 2458-2472.
- TARANTOLA, A. 1984, Inversion of seismic reflection data in the acoustic approximation, *Geophysics*, 49, 1259-1267.
- TAYLOR, M. 1981, *Pseudodifferential Operators*, Princeton University Press, Princeton, NJ.
- TREVES, F. 1980, *Introduction to Pseudodifferential and Fourier Integral Operators*, 1 and 2, Plenum Press, NY.
- WOLF, E. 1969, *Three dimensional structure of the earth*, Cambridge University Press, Cambridge.