



ELSEVIER

be conservative, or even volume-

classes of maps

$$\mathbf{(AA)} \quad (h_1^{-1} \cdots h_m^{-1})t(h_m \cdots h_1)t$$

$$\mathbf{(EA)} \quad (h_1^{-1} \cdots h_m^{-1})e_{m+1}(h_m \cdots h_1)t$$

$$\mathbf{(EE)} \quad (th_1^{-1} \cdots h_m^{-1})e_{m+1}(h_m \cdots h_1t)e_0$$

where h_i represents a Hénon transformation in the form (2) a

Theorem 2 (cf. [9, Corollary 2.3] or [15, Theorem 4.4]). *Two reduced words $g_m \cdots g_1$ and $g_n \cdots g_1$ represent the same polynomial automorphism g if and only if $n = m$ and there exist maps $s_i \in \mathcal{S}_1$, $i = 0, \dots, m$ such that $s_0 = s_m = \text{id}$ and $g_i = s_i g_i s_{i-1}^{-1}$.*

From this theorem it follows that

To prove the second part of the proposition, consider first a linear, nonelementary involution $a(x, y)$. In that case, taking $s(x, y) = x(1, 0) + ya(1, 0)$, we see that $a = sts^{-1}$.

Next, we show that every affine, nonelementary involution (12) is \mathfrak{g} -conjugate to its linear part a . We know that $(\cdot, \cdot) = (a - \text{id})(c, 0)$ for some scalar c . Taking $s(x, y) = (x + c, y)$ it follows that $sas^{-1} = a$ and the proof is complete. \square

3.2. Normal forms

We intend to d

A. Gómez, J.D. Me

Proof. Consider g given by the reduced word (14

